

Table 23. Preharvest Intervals (Days) and Re-Entry Intervals for Fungicides Registered for Use on Midwest Vegetables in 2008^a

	acibenzolar-S-methyl	azoxystrobin	boscalid	boscalid/pyraclostrobin	chlorothalonil	cyazofamid	cymoxanil/famoxadone	cyprodinil/fludioxonil	dimethomorph	fixed copper	Aluminum trisfosetyl-Al	iprodione	mancozeb	maneb	myclobutanil	phosphorous acid	propamocarb	propiconazole	pyraclostrobin	thiophanate methyl	trifloxystrobin	triflumizole	
Asparagus		100			190						110		180		180	0							
Bean, Lima		0	7		14			7		0		b		30		0				28			
Bean, Snap		0	7		7			7		0		b			0	0				14			
Beet		0								0												7	
Broccoli	7	0	0		7			7	0	0	3	0		7		0							
Brussels Sprouts	7	0	0		7			7	0	0	3			7									
Cabbage	7	0	0		7			7	0	0	3	0		7		0							
Cabbage, Chinese	7	0	c		7			7	0	0	3			7		0							
Cantaloupe		1		0	0	0	3		0	0	1/2		5	5	0	0	2		0	1	0	0	0
Carrot		0	0	0	0			7		0		0							0		7		
Cauliflower	7	0	0		7			7	0	0	3	0		7		0							
Celery		0			7			0		0	3							14				7	
Collard	7	0	14					7	0	0	3			14									
Cucumber		1		0	0	0	3		0	0	1/2		5	5	0		2		0	1	0	0	0
Eggplant		0	0							0				5		0			0		3		
Endive		0						0			3			10									
Kale	7	0	14					7			3			10									
Lettuce, Head		0	0	14			3	0	0		3	14		10			2						
Lettuce, Leaf		0	0	14			3	0	0		3	14		10			2						
Mint		7			80																		
Mustard		0	14					7		0	3												
Onion, Bulb		0	7	7	7			7	0	0	7	7	7	7		0			7				
Onion, Green		0	7	7	14			7	0	0				7		0			7				
Parsley		0						0		0	3												
Parsnip		0			10																	7	
Peas		0								0						0							
Pepper		0	0				3			0				7		0	5		0		3		
Potato		14	30	14	7	7	14		4	0		14	14	14		0	14		3	21	7		
Pumpkin		1		0	0	0	3		0	0	1/2		5	5	0		2		0	1	0	0	0
Radish		0																	0				
Spinach	7	0								0	3												
Squash, Summer		1		0	0	0	3		0	0	1/2		5	5	0		2		0	1	0	0	0
Squash, Winter		1		0	0	0	3		0	0	1/2		5	0		2		0	1	0	0	0	0
Sweet Corn		7			14								7	7				14					
Tomato	14	0	0		0	0	3		4	0	14		5	5	0	0	5		0		3		
Turnip	7	0								0												7	
Watermelon		1		0	0	0	3		0	0	1/2		5	5	0	0	2		0	1	0	0	0
Re-Entry Interval (hr)	12	4	12	12	12	12	12	12	12	24	12	24	24	24	24	4	12	24	12	12	12	12	12

^a Check label directions before applying any of these pesticides.
^b Do not apply past peak bloom.
^c 0-day PHI for Napa Chinese Cabbage, 14-day PHI for bok choy.